



MPFL Allograft Reconstruction Protocol Dr. Green

| | Treatment | Goals |
|-------------|--|--|
| Post-Op Day | -Dressing change/removal | -Independent in HEP |
| | -Instruct in signs/symptoms of infection | -Fair quad set |
| | -Check for DVT | -Independent with SLR |
| | -Cryotherapy | -Understand importance |
| | -Brace locked in extension | of knee immobilizer |
| | -NWB AROM to tolerance, progress slowly | |
| | -NWB x 6 weeks (unless otherwise stated in op report) | |
| | HEP : Heel slides, quad set, hamstring set, patellar mobs, gastroc | |
| | stretch, SLR in brace | |
| | -Modalities as needed | |
| Phase I | Frequency: TBD by PT upon assessment of quad function | -Continue HEP |
| 0-6 weeks | -NWB x 6 weeks | -Quad |
| | -Brace should be worn at all times during the day and while | strength/stabilization |
| | sleeping. Off for hygiene. | -Maintain WB restrictions |
| | -Progressing AROM to tolerance | |
| | -Continue table exercises, blood flow restriction (BFR) | |
| Phase II | -Progress to WBAT with t-scope locked in extension and then | -Maintain full extension |
| 6 weeks – 4 | into flexion per quad stability | -Full AROM |
| months | -Discontinue brace if able to achieve full extension without lag | -Normalize gait |
| | (Brace may be changed to patellar-stabilizing brace per MD orders) | |
| | Progress: | |
| | -WB flexibility, toe raises, closed-chain quad strength | |
| | -Floor based core and glute exercises, balance, hamstring curls, | |
| | core/pelvic, stability work and stationary biking; progress per | |
| | tolerance | |
| | -Full weight-bearing | |
| | -No bracing required unless otherwise ordered | |
| | -Begin elliptical, in-line jogging at 12 weeks under PT | |
| Phase III | supervision | -Full AROM |
| | Progress: | -i uli ANUIVI |
| 4-6 months | -Flexibility/strengthening, progression of function | Poturn to play as |
| | -Forward/backward running, cutting, grapevine | Return to play as tolerated after 16 weeks |
| | -Initiate plyometric program and sports specific drills at 16 weeks | |
| | weeks | post op when cleared by MD |